

280 S. Main Street Suite 200 Orange, CA 92868 + Tel. (714) 634-4567 + Fax (714) 634-4569 16300 Sand Canyon Ave · Suite 511 · Irvine, CA 92618 + Tel. (949) 255-9890 + Fax (949) 255-9776 Dr. Jeremy Smith + Direct Line Tel. (714) 937-2105 + Direct Fax (714) 937-4893

WHAT TO EXPECT WITH A TOTAL DISC REPLACEMENT

Please be advised that you must avoid all strenuous activity for 2 weeks after surgery or until your first post-operative visit.

LENGTH OF STAY

• Usually patients are discharged from the hospital the same day unless otherwise advised.

INCISIONS

• Usually vertical and 2-3 inches long.

MOBILITY

- The nursing and/or Physical Therapy staff will assist you with getting out of bed and walking around soon after surgery. There are no limitations to how much you can walk.
- PLEASE remember that although you have pain it is very important that you continue to walk through the pain. Walking will **greatly** decrease muscle spasm, pain and help with digestion.
- You will not begin physical therapy until your first post-operative visit.

COLLAR

- You will be given a soft collar to wear post operatively. This collar is to be worn at all times, except for showering, range of motion exercises, and eating, until we see you at your first post-operative appointment.
- Collar must be worn while sleeping.

MEDICATION

- You will likely receive a prescription at your pre-operative appointment for pain and muscle spasms. It is recommended that you take an over the counter stool softener for the first two weeks as well (ie. SennaS).
- Narcotics are very effective for pain relief but may cause other side effects. The possible effects vary among patients and may include: sleepiness, nausea, constipation, flushing, sweating, and occasionally euphoria or confused feelings.
- You will also be given a prescription for Toradol to be taken for 5 days post operatively. This will help the instrumentation integrate into the bone.

PAIN

- You may experience surgical discomfort after your procedure.
- You may experience "reminder" pain after your surgery. This is pain that is similar to your preoperative pain, sometimes on the opposite side, that will go away on its own with time. This pain is due to the postoperative swelling and irritation of the nerves.

OUESTIONS or CONCERNS

• If you have any additional questions/concerns please contact us at **714.937.2105** Monday-Friday 8:00 am to 5:00 pm or **714.634.4567** after hours and weekends.